

Title

Firstname Lastname

Affiliation

Abstract

—!!!—an abstract is required—!!!—

Keywords: —!!!—at least one keyword is required—!!!—.

Abstract

This paper shows the application of mortality projection techniques (Lee Carter) to the projection of retirement costs. The main purpose of the paper is to show how R can be successfully used to perform life expectancy projection with practical actuarial purpose for annuity business and social security insurance system. **demography** and **lifecontingencies** packages will be used. The analysis performed within this paper are mechanical and intended for didactic purposes.

1. Introduction

Mortality across any ages is showing continuous reduction across the world. In addition, a reliable projection of mortality is very important to accurately estimate the cost of life - insurance and pension annuities.

Methodologies have been developed to project the evolution of the mortality. Lee - Carter method is probably the most well known, among these.

The Lee Cartel forecasts results could be used to project a life table for each specific cohort (year) of birth on which pension annuities projection could be fit.

Our exercise will be based on Italian data downloaded from the Human Mortality Databases via **demography** package dedicated function. We will use **demography** and **forecast** package to fit Lee - Carter model and perform 100 - years in advance extrapolations.

Finally, **lifecontingencies** package will be used to project the cost of a pension annuity, $\ddot{a}_x^{(m)}$ for the cohorts of 1940, 1950, . . . , 2000 borns cohorts. Following demographic and economic assumptions will be hold:

- x , the retirement age will be set equal to 65 regardless the cohort.
- m , the number of fractional payments per year, will be equal to 12.
- $\ddot{a}_x^{(m)}$ to be the actuarial present value of a yearly annuity of 1 monetary unit, that will be evaluated assuming an interest rate of 4% and an inflation rate of 2%.

The projection has been performed using a mechanical approach, since the purpose of this paper lies in showing the procedure instead of providing sensible results. Finally, most of this paper is based on the examples provided in [Charpentier](#).

2. Fitting Lee Carter model

Lee Carter original model [Lee and Carter \(1992\)](#) assumes that the mortality (hazard rate) for age x in calendar year t can be expressed as Equation 1.

$$\ln \mu_{x,t} = a_x + b_x k_t + \varepsilon_{x,t} \quad (1)$$

Term in Equation 1 can be interpreted as follows:

1. a_x to be a base mortality rate for age x .
2. k_t to be the calendar year - related latent variable factor.
3. b_x is the sensitivity of age x to factor k_t .
4. $\varepsilon_{x,t}$ represents the residual component.

The one - year survival probability at age x during calendar year t is expressed by Equation 2. A longitudinal life table for the cohort of born in calendar year YYYY can be created selecting all $p_{x,t}$ for which $t - x = YYYY$.

$$p_{x,t} = \exp(-\mu_{x,t}) \quad (2)$$

We will perform such exercise on Italy HMD data, saved in mortalityDataset db.


```
R> library(demography)
R> library(forecast)
R> library(lifecontingencies)
R> load(file="mortalityDatasets.RData")
```

Following code creates a demogdata object from HDM data structure.

```
R> italyDemo<-demogdata(data=italyMx$rate$total,
+ F                pop=italyMx$pop$total,
+ F                name="total",
+ F                ages=italyMx$age,
+ F                years=italyMx$year,
+ F                type="mortality",
+ F                label="Italy",lambda=1)
```

Plot method is available on demogdata.

```
R> plot(italyDemo)
```


Italy: total death rates (1872–2008)

To fit Lee - Carter model (without going through logarithms) we do

```
R> italyLca<-lca(italyDemo)
```

`lca` returned object allows us to inspect a_x , b_x and k_t .

```
R> par(mfrow=c(1,3))
R> plot(x=italyLca$age, y=italyLca$ax, main="ax")
R> plot(x=italyLca$age, y=italyLca$bx, main="bx")
R> plot(x=italyLca$year, y=italyLca$kt, main="kt")
```




We can therefore use **forecast** package to project the future k_t s (up to 110).

```
R> ktSeries<-italyLca$kt
R> ktArima<-auto.arima(ktSeries,allowdrift=TRUE,max.order=20)
R> ktArimaForecasts<-forecast(ktArima, h=110)
R> fullKt<-ts(c(ktArimaForecasts$fitted, ktArimaForecasts$mean),start=1872)
```

and project results

```
R> plot(fullKt)
```



Then we create a function to project life table depending by year of birth.

```
R> createActuarialTable<-function(yearOfBirth){
+ F      #get projected Px
+ F      ktSubset<-window(fullKt, start=yearOfBirth)
+ F      predictionTable<-data.frame(age=italyLca$age,ax=italyLca$ax,bx=italyLca$bx)
+ F      predictionTable$kt=ktSubset[1:nrow(predictionTable)]
+ F      predictionTable$mux=with(predictionTable,exp(ax+bx*kt))
+ F      predictionTable$px=with(predictionTable,exp(-mux))
+ F      fittedPx=predictionTable$px
+ F      px4Completion=seq(from=predictionTable$px[length(fittedPx)], to=0, length=20)
+ F      totalPx=c(fittedPx,px4Completion[2:length(px4Completion)])
+ F      #create life table
+ F      irate=1.04/1.02-1
+ F
+ F      cohortLt=probs2lifetable(probs=totalPx, radix=100000,type="px", name=paste("Cohor
+ F      cohortAct=new("actuarialtable",x=cohortLt@x, lx=cohortLt@lx,
+ F      interest=irate, name=cohortLt@name)
+ F      return(cohortAct)
+ F      }
R>
```

We can therefore calculate the APV of $\ddot{a}_{65}^{(12)}$ for the selected cohorts.

```
R>      getAnnuityAPV<-function(yearOfBirth) {
```

```

+ F          actuarialTable<-createActuarialTable(yearOfBirth)
+ F          out=axn(actuarialTable,x=65,m=12)
+ F          return(out)
+ F      }
R>      for(i in seq(1920,2000,by=10)) {
+ F          cat("For cohort ",i, " the expected lifetime at birth is",
+ F              round(exn(createActuarialTable(i)),2),
+ F              " and the APV is :",round(getAnnuityAPV(i),2),"\n")
+ F      }

```

For cohort 1920 the expected lifetime at birth is 51.75 and the APV is : 5.45
For cohort 1930 the expected lifetime at birth is 63.15 and the APV is : 6.35
For cohort 1940 the expected lifetime at birth is 67.77 and the APV is : 7.09
For cohort 1950 the expected lifetime at birth is 76.19 and the APV is : 7.71
For cohort 1960 the expected lifetime at birth is 80.6 and the APV is : 8.29
For cohort 1970 the expected lifetime at birth is 82.93 and the APV is : 8.84
For cohort 1980 the expected lifetime at birth is 85.68 and the APV is : 9.35
For cohort 1990 the expected lifetime at birth is 87.96 and the APV is : 9.82
For cohort 2000 the expected lifetime at birth is 89.51 and the APV is : 10.27

References

- Charpentier A (????). "Actuarial Science with R 2: life insurance and mortality tables." <http://freakonometrics.blog.free.fr/index.php?post/2012/04/04/Life-insurance,-with-R,-Meielisalp>. Accessed: 04/11/2012.
- Lee R, Carter L (1992). "Modeling and forecasting U.S. mortality." *Journal of the American Statistical Association*, **87**(419), 659–675. doi:10.2307/2290201.

Affiliation:

Giorgio Alfredo Spedicato
StatisticalAdvisor
Via Firenze 11 20037 Italy
Telephone: +39/334/6634384
E-mail: lifecontingencies@statisticaladvisor.com
URL: www.statisticaladvisor.com