

Package ‘neonSoilFlux’

November 22, 2025

Type Package

Title Compute Soil Carbon Fluxes for the National Ecological
Observatory Network Sites

Version 3.0.0

Description Acquires and synthesizes soil carbon fluxes at sites located in the National Ecological Observatory Network (NEON). Provides flux estimates and associated uncertainty as well as key environmental measurements (soil water, temperature, CO₂ concentration) that are used to compute soil fluxes.

URL <https://github.com/jmzobitz/neonSoilFlux>

BugReports <https://github.com/jmzobitz/neonSoilFlux/issues>

License GPL (>= 3)

Encoding UTF-8

LazyData true

RoxygenNote 7.3.2

Depends R (>= 4.1)

Imports dplyr, ggplot2, lubridate, neonUtilities, purrr, stats,
stringr, tibble, tidyr, tidyselect, rlang

Suggests knitr, rmarkdown

VignetteBuilder knitr

NeedsCompilation no

Author John Zobitz [aut, cre] (ORCID: <<https://orcid.org/0000-0002-1830-143X>>),
Edward Ayres [aut] (ORCID: <<https://orcid.org/0000-0003-1846-1473>>),
Katie O'Rourke [ctb] (LinkedIn,
<<https://www.linkedin.com/in/katie-o-rourke-5a898b194/>>),
Zoey Werbin [ctb],
Lajntxiag Lee [ctb],
Ridwan Abdi [ctb] (LinkedIn,
<<https://www.linkedin.com/in/ridwaan-cabdi/>>),
Dijonë Mehmeti [ctb] (LinkedIn,
<<https://www.linkedin.com/in/dijon%C3%AB-mehmeti/>>),
Ly Xiong [ctb] (LinkedIn,
<<https://www.linkedin.com/in/ly-xiong-bb83ba1b2/>>)

Maintainer John Zobitz <zobitz@augsborg.edu>

Repository CRAN

Date/Publication 2025-11-22 19:40:02 UTC

Contents

acquire_neon_data	3
check_qf_flags	4
co2_to_umol	4
compute_flux_gradient_layer	5
compute_monthly_mean	6
compute_neon_flux	7
compute_surface_flux	8
compute_surface_flux_layer	9
correct_env_data	9
dejong_shappert_flux	10
depth_interpolate	11
determine_position	12
diffusivity	12
env_fingerprint_plot	13
fit_function	14
flux_fingerprint_plot	15
hirano_flux	15
insert_mean	16
inside_interval	17
linear_regression	18
measurement_detect	19
measurement_merge	19
neonBoulded60Cal	20
neonCal	21
powerset	22
quadrature_error	23
reprocess_vswc	24
sjer_env_data_2022_06	24
sjer_flux_2022_06	25
sjer_megapit_data_2022_06	26
swc_correct	27
swc_corrections	28
tang_2003_flux	28
tang_2005_flux	29

Index	31
--------------	-----------

acquire_neon_data	<i>Acquire NEON data for processing</i>
-------------------	---

Description

Given a site code and dates, apply the neonUtilities package to download the data from NEON API

Usage

```
acquire_neon_data(  
  site_name,  
  download_date,  
  time_frequency = "30_minute",  
  provisional = FALSE  
)
```

Arguments

site_name	Required. NEON code for a particular site (a string)
download_date	Required. Date where we end getting NEON data. Format: YYYY-MM (can't specify day). So "2020-05" means it will grab data for the entire 5th month of 2020. (a string). Downloads data for a given month only
time_frequency	Required. Will you be using 30 minute ("30_minute") or 1 minute ("1_minute") recorded data? Defaults to 30 minutes.
provisional	Required. Should you use provisional data when downloading? Defaults to FALSE. See NEON Data Releases . Defaults to FALSE (similar to include.provisional in loadByProduct).

Value

A list containing stacked environmental data ('site_data') and soil properties ('site_megapit').

Author(s)

John Zobitz <zobitz@augsborg.edu>

Examples

```
out_env_data <- acquire_neon_data("SJER", "2022-06")
```

check_qf_flags	<i>Internal helper function to determine availability of data within a time interval</i>
----------------	--

Description

Given a data measurement, determine when a monthly mean is used and the bulk QF flags. Helps to determine when the environmental measurements produced a QF value and to be used in subsequent flux calculations.

Usage

```
check_qf_flags(measurement_name, data)
```

Arguments

measurement_name	The name of the measurement (staPres, soilTemp, VSWC, soilCO2Concentration)
data	Data used to check the qf flags

Value

A data frame of startDateTime, horizontalPosition, and the associated QF flag.
 # changelog and author contributions / copyrights

Author(s)

John Zobitz <zobitz@augzburg.edu>

co2_to_umol	<i>Convert co2 concentration from ppm to $\mu\text{mol m}^{-3}$ units</i>
-------------	--

Description

Given a measurement of co2, convert it from ppm to $\mu\text{mol m}^{-3}$ based on temperature and pressure. Also compute associated error via quadrature.

Usage

```
co2_to_umol(
  temperature,
  pressure,
  co2,
  temperature_err,
  pressure_err,
  co2_err,
  zoffset
)
```

Arguments

temperature	Required. Soil temperature (degrees C)
pressure	Required. Barometric air pressure (kilopascal)
co2	Carbon dioxide in ppm
temperature_err	Required. Reported Soil temperature error (degrees C)
pressure_err	Required. Reported Barometric air pressure error (kilopascal)
co2_err	Required. Carbon dioxide in ppm error
zoffset	Required. Surface depth (m). Reported as a negative number.

Value

A value of the converted co2

Author(s)

John Zobitz <zobitz@augsborg.edu>

Examples

```
co2_to_umol(31,96.3,654,.15,.05,9,-.05)
```

```
compute_flux_gradient_layer
```

Computes Fick's law of diffusion between two values

Description

Computes Fick's law of diffusion between two values

Usage

```
compute_flux_gradient_layer(depths, co2, co2_err, diffus, diffus_err)
```

Arguments

depths	Required. Depths that the compute the flux (2 values)
co2	Required. CO2 molar concentration (2 values, corresponding to depths)
co2_err	Required. Error in co2 (2 values, corresponding to depths)
diffus	Required. Diffusivity (a single value)
diffus_err	Required. Diffusive error (a single value)

Value

Tibble of depth, flux, and error. The depth is the mean between the depths

Examples

```
compute_flux_gradient_layer(depths = c(-0.03,-0.09),
  co2 = c(28030, 66834.66),
  co2_err = c(1327,1246),
  diffus = 7.13e-06,
  diffus_err = 3.44e-6
)
```

compute_monthly_mean *Function to compute monthly means for a given month of NEON data.*

Description

Given a NEON measurement data frame calculate the monthly mean values across all horizontal and vertical locations. Based off code from Zoey Werbin.

Usage

```
compute_monthly_mean(
  NEON_data,
  time_horizon = 15,
  position_columns = c("horizontalPosition", "verticalPosition")
)
```

Arguments

NEON_data	Required. Input vector of neon measurements for a month
time_horizon	Required. How many unique days do we need each month?
position_columns	Optional. Do we group by horizontalPosition, verticalPosition, and? Default is both. Added this option in case we just want to average across a given dimension.

Value

A data frame that reports for each horizontal and vertical position the computed mean and standard deviation from sampling (similar to a bootstrap method) as well as the sample mean and sample standard deviation

Author(s)

John Zobitz <zobitz@augsborg.edu>

References

Zoey Werbin (@zoey-rw): original author <https://github.com/zoey-rw/microbialForecasts/blob/caa7b1a8aa8a131a5ff9340f15>

Examples

```
# Download the NEON data directly - here this would be soil moisture
NEON_moist_30m_orig <- neonUtilities::loadByProduct(
  dpID = "DP1.00094.001",
  site = "WREF",
  startdate = "2022-06",
  enddate = "2022-06",
  timeIndex = "30",
  package = "expanded",
  check.size = FALSE,
  include.provisional = TRUE
)

# Then correct the swc
site_swc <- swc_correct(NEON_moist_30m_orig, "WREF", "2022-06")

# Select the columns and the time frequency
time_frequency <- "30_minute"
column_selectors <- c("Mean", "Minimum", "Maximum", "ExpUncert", "StdErMean")

  swc <- site_swc |>
  purrr::pluck(paste0("SWS_", time_frequency)) |>
  dplyr::select(tidyselect::all_of(c("domainID", "siteID",
    "horizontalPosition", "verticalPosition", "startDateTime", "VSWCFinalQF")),
    tidyselect::matches(stringr::str_c("VSWC", column_selectors)))

# Determine a data frame of the different horizontal and vertical positions
swc_positions <- site_swc |>
  purrr::pluck(paste0("sensor_positions_", "00094"))

# Add on the positions for swc
swc <- determine_position(swc_positions, swc)
```

compute_neon_flux	<i>Compute NEON fluxes at a site</i>
-------------------	--------------------------------------

Description

Given a site filename (from `acquire_neon_data`), process and compute fluxes. This file takes a saved data file from `acquire`: 1) Takes the needed components (QF and measurement flags) for soil water, temperature, co2, binding them together in a tidy data frame 2) Interpolates across the measurements 3) Merges air pressure data into this data frame 4) Does a final QF check so we should have only timeperiods where all measurements exist 5) Adds in the megapit data so we have bulk density, porosity measurements at the interpolated depth. 6) Saves the data

Usage

```
compute_neon_flux(input_site_env, input_site_megapit)
```

Arguments

input_site_env Required. Input list of environmental data. Usually given from acquire_neon_data

input_site_megapit

Required. Input list of environmental soil data. Usually given from acquire_neon_data

Value

Data frame of fluxes and gradient from the timeperiod

Author(s)

John Zobitz <zobitz@augsborg.edu> based on code developed by Edward Ayres <eayres@battelleecology.org>

Examples

```
# First acquire the NEON data at a given NEON site
out_env_data <- acquire_neon_data("SJER", "2020-05")

# Then process and compute the fluxes:
out_flux <- compute_neon_flux(input_site_env = sjer_env_data_2022_06,
input_site_megapit = sjer_megapit_data_2022_06)
```

compute_surface_flux *Internal function to compute CO2 surface flux*

Description

Computation function. Given a measurement of the co2 and diffusive flux at different levels, return the surface flux

Usage

```
compute_surface_flux(input_data)
```

Arguments

input_data Required. A data frame containing zOffsets, diffusivity, and co2 (umol mol-1) and their associated errors

Value

A value of the surface CO2 flux (umol m-2 s-1)

Author(s)

John Zobitz <zobitz@augsborg.edu>

`compute_surface_flux_layer`*Title Determine the surface flux from linear regression at layers*

Description

Computation function. Given a measurement of the co2 and diffusive flux at different levels, return the surface flux

Usage

```
compute_surface_flux_layer(input_data)
```

Arguments

<code>input_data</code>	Required. A data frame containing zOffsets, diffusivity, and co2 (umol mol-1) and their associated errors
-------------------------	---

Value

A value of the surface CO2 flux (umol m-2 s-1)

Author(s)

John Zobitz <zobitz@augsborg.edu>

See Also

[`compute_surface_flux()`] for other ways to compute surface fluxes.

`correct_env_data`*Internal function that prepares downloaded NEON data for flux processing*

Description

This file takes data frame from `acquire_neon_data` and: 1) Takes the needed components (QF and measurement flags) for soil water, temperature, co2, binding them together in a tidy data frame 2) Interpolates across the measurements 3) Merges air pressure data into this data frame

Usage

```
correct_env_data(input_data)
```

Arguments

<code>input_data</code>	Required. Nested data frame from <code>acquire_neon_data</code> .
-------------------------	---

Value

List of all QF flags over time period and Data frame of environmental measurements for flux computation

Author(s)

John Zobitz <zobitz@augzburg.edu>

Examples

```
# Note: you may need to first acquire the NEON data using acquire_neon_data
# Now correct existing environmental data:
corrected_data <- correct_env_data(sjer_env_data_2022_06)
```

dejong_shappert_flux *Internal function to compute surface co2 flux at a given timepoint via De Jong and Schappert (1972)*

Description

Given zOffsets, diffusivity, and co2 (umol mol⁻¹) and their associated errors, compute the surface flux. This is done by estimating the surface concentration (through linear regression) and doing a gradient calculation. Modified from De Jong and Schappert (1972).

Usage

```
dejong_shappert_flux(zOffset, co2, co2_err, diffusive, diffusive_err)
```

Arguments

zOffset	Required. depths below surface - assumed to be positive in value. Important for directionality!
co2	Required. co2 at depth (umol m ⁻¹)
co2_err	Required. Associated errors with that value of co2
diffusive	Required. diffusivity at each depth
diffusive_err	Required Associated errors with diffusivity

Value

Data frame of fluxes associated error

Author(s)

John Zobitz <zobitz@augzburg.edu>

References

- Jong, E. De, and H. J. V. Schappert. 1972. "Calculation of Soil Respiration and Activity from CO₂ Profiles in the Soil." *Soil Science* 113 (5): 328.
- Maier, M., and H. Schack-Kirchner. 2014. "Using the Gradient Method to Determine Soil Gas Flux: A Review." *Agricultural and Forest Meteorology* 192–193 (July):78–95. <https://doi.org/10.1016/j.agrformet.2014.03.006>.

See Also

[hirano_flux()], [tang_2003_flux()], [tang_2005_flux()] for other ways to compute surface fluxes.

depth_interpolate	<i>Internal function to interpolate different depth measurements</i>
-------------------	--

Description

Definition function. Linearly interpolate a measurement across the different measurement depths

Usage

```
depth_interpolate(
  input_measurements,
  measurement_name,
  measurement_interpolate
)
```

Arguments

input_measurements	Required. Nested data frame (tibble of a months worth of data of co2, temperature, swc, pressure)
measurement_name	Required. Names of measurements we are interpolating. Currently only does one column at a time.
measurement_interpolate	Required. Names of measurement whose depth is used to interpolate (typically co2)

Value

A nested data frame with interpolated measurements.

Author(s)

John Zobitz <zobitz@augsborg.edu> based on code developed by Edward Ayres <eayres@battelleecology.org>

determine_position	<i>Internal function to determine the depth of a measurement</i>
--------------------	--

Description

Given a NEON measurement data frame and measurement depth, pull out the measurement depth for a measurement - because it may vary in time based on if a sensor is replaced.

Usage

```
determine_position(input_positions, input_measurement)
```

Arguments

input_positions	Required. Input vector of measurements depths
input_measurement	Required. Input vector of measurements.

Value

A data frame that reports the measurement depth for and associated environmental measurement.

Author(s)

John Zobitz <zobitz@augsborg.edu>

diffusivity	<i>Compute soil diffusivity</i>
-------------	---------------------------------

Description

Given a tidied data frame of soil measurements (from interpolate.R), compute the diffusivity in a given soil layer

Usage

```
diffusivity(
  temperature,
  soil_water,
  pressure,
  temperature_err,
  soil_water_err,
  pressure_err,
  zOffset,
  porVol2To20
)
```

Arguments

temperature	Required. Soil temperature (degrees C)
soil_water	Required. Soil water content
pressure	Required. Barometric air pressure (kilopascal)
temperature_err	Required. Reported Soil temperature error (degrees C)
soil_water_err	Required. Reported Soil water content error
pressure_err	Required. Reported Barometric air pressure error (kilopascal)
zOffset	Required. Measurement level in cm.
porVol2To20	Required. Porosity of the 0-20 mm fraction (cm ³ cm ⁻³). Assumes no pores within rocks.

Value

A value of the computed diffusivity

Author(s)

John Zobitz <zobitz@augzburg.edu>

Examples

```
diffusivity(31,0.0102,96.3,.15,.2135,.05,-.05,0.45)
```

env_fingerprint_plot *Helper function to plot QF results for environmental measurements.*

Description

Given a flux measurement data frame, show when the environmental measurements produced a QF value

Usage

```
env_fingerprint_plot(input_fluxes)
```

Arguments

input_fluxes data frame of computed fluxes

Value

A ggplot graph where we have ordered factors showing the QA values a given environmental measurement

Author(s)

John Zobitz <zobitz@augzburg.edu>

Examples

```
# Make a fingerprint plot for environmental variables:
env_fingerprint_plot(sjer_flux_2022_06)
```

fit_function	<i>Internal function that interpolates a soil measurement to different depths</i>
--------------	---

Description

Definition function. Linearly interpolate a measurement across the different measurement depths

Usage

```
fit_function(
  input_depth,
  input_value,
  input_value_err,
  input_value_qf,
  interp_depth,
  measurement_special
)
```

Arguments

input_depth	Required. Vector of measurement depths.
input_value	Required. Vector of measured values (i.e. soil temperature and soil water) measured at depths input_depth
input_value_err	Required. Vector or reported measurement errors. Used to compute prediction error (via quadrature) when linear interpolation is used.
input_value_qf	Required. Vector of qf values from the smoothing 0 = no smoothing, 1 = mean value used (smoothing), 2 = NA value
interp_depth	Depths of the sensors required for interpolation
measurement_special	Flag if we want to just do linear interpolation for a given measurement

Value

A data frame of the depth and the measured column for the measurements and reported error

Author(s)

John Zobitz <zobitz@augzburg.edu> based on code developed by Edward Ayres <eayres@battelleecology.org>

flux_fingerprint_plot *Helper function to plot QF results for fluxes.*

Description

Given a flux measurement data frame, show when the flux and diffusivity measurements produced a QF value

Usage

```
flux_fingerprint_plot(input_fluxes, input_diffus_method = "Marshall")
```

Arguments

input_fluxes data frame of computed fluxes
input_diffus_method character string of diffusivity method used to compute fluxes. Can be one of "Marshall" or "Millington-Quirk".

Value

A ggplot graph where we have ordered factors showing the QA values a given flux computation

Author(s)

John Zobitz <zobitz@augsborg.edu>

Examples

```
# Make a fingerprint plot for computed flux values:
flux_fingerprint_plot(sjer_flux_2022_06)
```

hirano_flux	<i>Internal function to compute surface co2 flux at a given timepoint via Hirano et al 2005</i>
-------------	---

Description

Given zOffsets, diffusivity, and co2 (umol mol⁻¹) and their associated errors, compute the surface flux. This is done by estimating the surface concentration through linear regression and linear extrapolation of the bottom and top surface fluxes. Modified from Hirano et al (2005).

Usage

```
hirano_flux(zoffset, co2, co2_err, diffusive, diffusive_err)
```

Arguments

zOffset	Required. depths below surface - assumed to be positive in value. Important for directionality!
co2	Required. co2 at depth (umol m ⁻¹)
co2_err	Required. Associated errors with that value of co2
diffusive	Required. diffusivity at each depth
diffusive_err	Required Associated errors with diffusivity

Value

Data frame of fluxes associated error

Author(s)

John Zobitz <zobitz@augzburg.edu>

References

Hirano, Takashi, Honghyun Kim, and Yumiko Tanaka. 2003. "Long-Term Half-Hourly Measurement of Soil CO₂ Concentration and Soil Respiration in a Temperate Deciduous Forest." *Journal of Geophysical Research: Atmospheres* 108 (D20). <https://doi.org/10.1029/2003JD003766>.

Maier, M., and H. Schack-Kirchner. 2014. "Using the Gradient Method to Determine Soil Gas Flux: A Review." *Agricultural and Forest Meteorology* 192–193 (July):78–95. <https://doi.org/10.1016/j.agrformet.2014.03.006>.

See Also

[dejong_shappert_flux()], [tang_2003_flux()], [tang_2005_flux()] for other ways to compute surface fluxes.

insert_mean	<i>Internal function that inserts smoothed mean value of a measurement at a site</i>
-------------	--

Description

Given a site measurement and monthly mean, insert in the monthly mean value when the QF flag fails.

Usage

```
insert_mean(data, monthly_mean, measurement_name)
```

Arguments

data	Required. input data to use
monthly_mean	Required. monthly mean of input data to use
measurement_name	Required. name of measurement

Value

Nested data frame of measurements

Author(s)

John Zobitz <zobitz@augzburg.edu>

inside_interval	<i>Determine if a YYYY-MM string is inside a interval</i>
-----------------	---

Description

Determine if a YYYY-MM string is inside a interval

Usage

```
inside_interval(start, end, reference_time)
```

Arguments

start	starting interval time
end	ending interval time
reference_time	time we are comparing to - YYYY-MM string

Value

Logical indicating whether or not the reference time is inside the interval. We need this when working with downloaded NEON data which usually comes in a YYYY-MM string

Author(s)

John Zobitz <zobitz@augzburg.edu>

Examples

```
# Define starting and ending dates:
start_date <- as.POSIXct("2021-06-01 09:30:00",tz="UTC")
end_date <- as.POSIXct("2023-06-01 20:00:00",tz="UTC")

# Test, should return TRUE
inside_interval(start_date,end_date,"2022-06")

# Test, should return FALSE
inside_interval(start_date,end_date,"2020-06")
```

linear_regression	<i>Compute linear regression coefficients and errors.</i>
-------------------	---

Description

Given x and y data compute simple linear regression slope and intercept, with errors by quadrature.

Usage

```
linear_regression(x, x_err, y, y_err)
```

Arguments

x	Required. vector of independent variables.
x_err	Required. vector of independent variable errors.
y	Required. Vector of dependent variables.
y_err	Required. Vector of dependent variable errors.

Value

A data frame of linear regression slope, intercept, and associated errors

Author(s)

John Zobitz <zobitz@augzburg.edu>

Examples

```
y<- c(345,432,233)
y_err <- c(1,5,6)
x <- c(0.01,0.04,0.17)
x_err <- c(0.1,.3,.6)
linear_regression(x,x_err,y,y_err)
```

measurement_detect	<i>Internal function that makes sure for each time, position, and depth we have at least two data points for soil temp and soil h20, 3 for soil co2</i>
--------------------	---

Description

Given a merged data frame of co2, water, and temperature: 1) Filters on QF measurement flags 2) Filters if we have at least 2 soil h20 and temperature measurements, 3 co2 measurements at each time, horizontal position, and vertical depth 3) Filters if we have at least 3 distinct measurements at each time and horizontal position 4) Returns the resulting data frame.

This internal function is created to speed up data processing.

Usage

```
measurement_detect(input_data)
```

Arguments

input_data	Required. Nested data frame of merged soil water, temperature, co2, and pressure across different NEON locations and depths
------------	---

Value

Data frame of fluxes from the timeperiod

Author(s)

John Zobitz <zobitz@augsborg.edu> based on code developed by Edward Ayres <eayres@battelleecology.org>

measurement_merge	<i>Internal function that filters environmental data for easier processing of fluxes.</i>
-------------------	---

Description

Internal function. Given a set of environmental measurements, create a merged data frame of measurements and positions where the QF_flag exists (either observed or a smoothed mean measurement).

Usage

```
measurement_merge(
  neon_data,
  data_code,
  data_product_id,
  measurement_name,
  qf_name
)
```

Arguments

neon_data	Required. A list of NEON data downloaded from the utilities
data_code	Required. Names of data product we are interpolating. (SWS = soil water, ST = soil temperature, SCO2C = soil CO2)
data_product_id	Name of the data product 00094 = soil water, 00041 = soil temperature, 00095 = soil CO2
measurement_name	Required. Names of column we are grabbing. (VSWCMean = soil water, soil-TempMean = soil temperature, soilCO2concentrationMean = soil CO2)
qf_name	Required. Names of qf column we are grabbing. (VSWCFinalQF = soil water, finalQF = soil temperature, finalQF = soil CO2)

Value

A data frame of the requested data.

Author(s)

John Zobitz <zobitz@augsborg.edu>

neonBoulded60Cal	<i>Soil water content calibrations for belowground sensor data</i>
------------------	--

Description

A datatble of SWC corrections to remove step changes in soil water content measureing depths

Usage

```
neonBoulded60Cal
```

Format

A data table with 1881 observations and 18 variables

Details

- domain: NEON sampling domain
- site: NEON site
- plot: Which particular NEON plot at the site
- measurementLevel: Which particular measurement depth at that plot
- sensorDepths: Corrected depth of sensor measurement
- CVALA1: calibration coefficient a
- CVALA2: calibration coefficient b

- CVALA3: calibration coefficient c
- CVALR1: revision control
- CVALR2: failed file
- U_CVALA1: soil moisture combined uncertainty
- U_CVALA2: soil moisture repeatability
- U_CVALA3: soil moisture combined uncertainty (truth and trueness only)
- CVALO1: coefficients manually overridden (Y/N)
- CVALS1: start date for the soil specific calibration
- CVALS2: end date for the soil specific calibration, NA if no end date is defined
- remarks: any comments about the soil specific calibration information
- CVALC1: information build date

Source

<<https://data.neonscience.org/data-products/DP1.00094.001>> and <<https://doi.org/10.6084/m9.figshare.24424762.v1>>

neonCal

Soil water content calibrations for belowground sensor data

Description

A datatble of SWC corrections to remove step changes in soil water content measureing depths

Usage

neonCal

Format

A data table with 1881 observations and 18 variables

Details

- domain: NEON sampling domain
- site: NEON site
- plot: Which particular NEON plot at the site
- measurementLevel: Which particular measurement depth at that plot
- sensorDepths: Corrected depth of sensor measurement
- CVALA1: calibration coefficient a
- CVALA2: calibration coefficient b
- CVALA3: calibration coefficient c
- CVALR1: revision control

- CVALR2: failed file
- U_CVALA1: soil moisture combined uncertainty
- U_CVALA2: soil moisture repeatability
- U_CVALA3: soil moisture combined uncertainty (truth and trueness only)
- CVALO1: coefficients manually overridden (Y/N)
- CVALS1: start date for the soil specific calibration
- CVALS2: end date for the soil specific calibration, NA if no end date is defined
- remarks: any comments about the soil specific calibration information
- CVALC1: information build date

Source

<<https://data.neonscience.org/data-products/DP1.00094.001>> and <<https://doi.org/10.6084/m9.figshare.24424762.v1>>

powerset	<i>Compute all possible combinations of a vector - ignoring the empty set</i>
----------	---

Description

Compute all possible combinations of a vector - ignoring the empty set

Usage

```
powerset(x, size_start = 1)
```

Arguments

x	Required. Input vector
size_start	First power set input size

Value

A list of possible options

Examples

```
# All possible combinations
powerset(1:4)

# Now try without ignoring the single sets:
powerset(1:4,2)
```

quadrature_error	<i>Helper function to quickly compute the quadrature error</i>
------------------	--

Description

Assume a derived quantity y is a function of inputs x_i : $y = f(x_1, x_2, x_3, \dots)$

Given uncertainties (x_{err}) for each x_i , then this function will compute the corresponding y_{err} via quadrature. Inputs are the vector of partial derivatives df/dx_i , evaluated at $(x_1, x_2, x_3, \dots)$.

Resulting y_{err} is the square root of the sum of $(df/dx_1)^2 * (x_{\text{err}})^2 + (df/dx_2)^2 * (x_{\text{err}})^2 + (df/dx_3)^2 * (x_{\text{err}})^2 \dots$

Usage

```
quadrature_error(x_pd, x_err)
```

Arguments

<code>x_pd</code>	Required. Input vector of partial derivatives for $y = f(x)$, evaluated at x_i
<code>x_err</code>	Required. Error vector of measurements

Value

A value of quadrature error

Author(s)

John Zobitz <zobitz@augsborg.edu>

Examples

```
# Let's say we have 5 temperature measurements w/ error::
temperature <- c(31.108, 30.689, 30.463, 30.381, 30.250)
temperature_error <- c(0.1508, 0.1507, 0.1497, 0.1496, 0.1497)

# The sample mean is the sum of all measurements divided by the average:
sum(temperature)/5 # (Can also be computed with mean(temperature))

# The vector of partial derivatives is just 1/n for each measurement:
temperature_pd <- c(1/5, 1/5, 1/5, 1/5, 1/5)
quadrature_error(temperature_pd, temperature_error)
# Note: quadrature_error(1/5, temperature_error) is also allowed.
```

reprocess_vswc	<i>Internal helper function to reprocess SWC data from calibration data</i>
----------------	---

Description

Apply correct calibration data functions for SWC data - see <https://data.neonscience.org/data-products/DP1.00094.001>

Usage

```
reprocess_vswc(site_name, download_date)
```

Arguments

site_name	Required. NEON code for a particular site (a string)
download_date	Required. Date where we end getting NEON data. Format: YYYY-MM (can't specify day). So "2020-05" means it will grab data for the entire 5th month of 2020. (a string). Downloads data for a given month only

Value

A data frame of startDateTime, horizontalPosition, and the associated QF flag.

A list containing stacked swc data

Author(s)

John Zobitz <zobitz@augzburg.edu>,

Examples

```
out_swc <- reprocess_vswc("SJER", "2022-06")
```

sjer_env_data_2022_06	<i>Measured environmental data at a NEON site</i>
-----------------------	---

Description

A nested dataset containing environmental variables at the SJER site from June 2022. A convenience data frame to make processing fluxes easier for testing. Computed with the function `acquire_neon_data`. All the variables can be used to compute fluxes (along with megapit data).

Usage

```
data(sjer_env_data_2022_06)
```

Format

A nested 3 item list

Details

- measurement: Name of environmental measurement
- data: Nested list of data corresponding to a measurement
- monthly_mean: Monthly mean of measurement, computed in ‘compute_monthly_mean’

sjer_flux_2022_06	<i>Computed flux values at a NEON site</i>
-------------------	--

Description

A nested dataset containing computed fluxes at the SJER site from June 2022. Fluxes were computed with using `compute_neon_flux(sjer_env_data_2022_06,sjer_megapit_data_2022_06)`.

Usage

```
data(sjer_flux_2022_06)
```

Format

A nested data frame item list with 7200 rows and 8 columns

Details

- startDateTime: Time period of measurement (as POSIXct)
- horizontalPosition: Sensor location where flux is computed
- flux_compute: A nested tibble with variables (1) flux, flux_err, and method (one of 4 implemented)
- diffusivity: Computation of surface diffusivity
- VSWCMeanQF: QF flag for soil water content across all vertical depths at the given horizontal position: 0 = no issues, 1 = monthly mean used in measurement, 2 = QF fail
- soilTempMeanQF: QF flag for soil temperature across all vertical depths at the given horizontal position: 0 = no issues, 1 = monthly mean used in measurement, 2 = QF fail
- soilCO2concentrationMeanQF: QF flag for soil CO2 concentration across all vertical depths at the given horizontal position: 0 = no issues, 1 = monthly mean used in measurement, 2 = QF fail
- staPresMeanQF: QF flag for atmospheric pressure across all vertical depths at the given horizontal position: 0 = no issues, 1 = monthly mean used in measurement, 2 = QF fail

`sjer_megapit_data_2022_06`*Measured soil physical properties at a NEON site*

Description

A nested dataset containing soil data properties at the SJER site from June 2022. A convenience data frame to make processing fluxes easier for testing. Computed with using the function `acquire_neon_data`. Essentially what is returned when data product DP1.00096.001 is loaded from [loadByProduct](#).

Usage

```
data(sjer_megapit_data_2022_06)
```

Format

A nested 10 item list

Details

- `citation_00096_RELEASE-2024`: BibTex reference
- `issueLog_00096`: Listing of known issues
- `mgp_perarchivesample`: Archived sample information
- `mgp_perbiogeosample`: Data collected on biogeochemistry sample
- `mgp_perbulksample`: Data collected on bulk density sample
- `mgp_perhorizon`: Per soil horizon metadata
- `mgp_permegapit`: Data collected per megapit
- `readme_00096`: Data product description, issue log, and other metadata about the data product
- `validation_00096`: Description of data validation applied at the points of collection and ingest
- `variables_00096`: Description and units for each column of data in data tables

Source

From <https://data.neonscience.org/data-products/DP1.00096.001>

swc_correct

*Internal function to correct depths for VSWC NEON data.***Description**

Given the expanded SWC data, return a corrected version based on the values below

Usage

```
swc_correct(input_swc, curr_site, reference_time)
```

Arguments

input_swc	Required. input soil water content data from <code>acquire_neon_data</code> (as a list)
curr_site	Current site we are working with
reference_time	Current month we are working with

Value

A revised list of corrected soil water content and depths.

Author(s)

John Zobitz <zobitz@augsborg.edu>

Examples

```
# Download the soil water content data:
site_swc <- neonUtilities::loadByProduct(
  dpID="DP1.00094.001",
  site="SJER",
  startdate="2020-05",
  enddate="2020-05",
  timeIndex = "30",
  package="expanded",
  check.size = FALSE,
  include.provisional = TRUE
)

# Then correct the swc:
site_swc <- swc_correct(site_swc,"SJER","2020-05")
```

swc_corrections	<i>Corrected sensor locations for NEON soil water content data</i>
-----------------	--

Description

A dataset containing corrected NEON sensor location depths for soil water content data

Usage

```
swc_corrections
```

Format

A data frame with 2161 rows and 6 variables

Details

- domainID (ecological domain of site)
- siteID NEON code to refer to site
- HOR.VER 3 digit code to refer to the horizontal and vertical position of measuring location.
- sensorDepth depth below surface (m)
- startDateTime time measurement was started
- endDateTime time measurement ended

Source

<<https://data.neonscience.org/data-products/DP1.00094.001>>

tang_2003_flux	<i>Internal function to compute surface co2 flux at a given timepoint via Tang et al 2003</i>
----------------	---

Description

Given zOffsets, diffusivity, and co2 (umol mol⁻¹) and their associated errors, compute the surface flux. This is done by computing the gradient of co2 (slope from linear regression) and linear extrapolation of surface diffusivity. Modified from Tang et al (2003).

Usage

```
tang_2003_flux(zOffset, co2, co2_err, diffusive, diffusive_err)
```

Arguments

zoffset	Required. depths below surface - assumed to be positive in value. Important for directionality!
co2	Required. co2 at depth (umol m ⁻¹)
co2_err	Required. Associated errors with that value of co2
diffusive	Required. diffusivity at each depth
diffusive_err	Required Associated errors with diffusivity

Value

Data frame of fluxes associated error

Author(s)

John Zobitz <zobitz@augzburg.edu>

References

Tang, Jianwu, Dennis D Baldocchi, Ye Qi, and Liukang Xu. 2003. "Assessing Soil CO₂ Efflux Using Continuous Measurements of CO₂ Profiles in Soils with Small Solid-State Sensors." *Agricultural and Forest Meteorology* 118 (3): 207–20. [https://doi.org/10.1016/S0168-1923\(03\)00112-6](https://doi.org/10.1016/S0168-1923(03)00112-6).

Maier, M., and H. Schack-Kirchner. 2014. "Using the Gradient Method to Determine Soil Gas Flux: A Review." *Agricultural and Forest Meteorology* 192–193 (July):78–95. <https://doi.org/10.1016/j.agrformet.2014.03.006>.

See Also

[dejong_shappert_flux()], [hirano_flux()], [tang_2005_flux()] for other ways to compute surface fluxes.

tang_2005_flux	<i>Internal function to compute surface co2 flux at a given timepoint via Tang et al 2005</i>
----------------	---

Description

Given zOffsets, diffusivity, and co2 (umol mol⁻¹) and their associated errors, compute the surface flux. This is done by linear extrapolation of surface fluxes. Modified from Tang et al (2005).

Usage

```
tang_2005_flux(zoffset, co2, co2_err, diffusive, diffusive_err)
```

Arguments

zOffset	Required. depths below surface - assumed to be positive in value. Important for directionality!
co2	Required. co2 at depth (umol m ⁻¹)
co2_err	Required. Associated errors with that value of co2
diffusive	Required. diffusivity at each depth
diffusive_err	Required Associated errors with diffusivity

Value

Data frame of fluxes associated error

Author(s)

John Zobitz <zobitz@augsborg.edu>

References

- Tang, Jianwu, Laurent Misson, Alexander Gershenson, Weixin Cheng, and Allen H. Goldstein. 2005. "Continuous Measurements of Soil Respiration with and without Roots in a Ponderosa Pine Plantation in the Sierra Nevada Mountains." *Agricultural and Forest Meteorology* 132 (3): 212–27. <https://doi.org/10.1016/j.agrformet.2005.07.011>.
- Maier, M., and H. Schack-Kirchner. 2014. "Using the Gradient Method to Determine Soil Gas Flux: A Review." *Agricultural and Forest Meteorology* 192–193 (July):78–95. <https://doi.org/10.1016/j.agrformet.2014.03.006>.

See Also

[dejong_shappert_flux()], [hirano_flux()], [tang_2003_flux()] for other ways to compute surface fluxes.

Index

- * **Currently**
 - compute_monthly_mean, [6](#)
- * **datasets**
 - neonBoulded60Cal, [20](#)
 - neonCal, [21](#)
 - sjer_env_data_2022_06, [24](#)
 - sjer_flux_2022_06, [25](#)
 - sjer_megapit_data_2022_06, [26](#)
 - swc_corrections, [28](#)
- * **none**
 - compute_monthly_mean, [6](#)
- acquire_neon_data, [3](#)
- check_qf_flags, [4](#)
- co2_to_umol, [4](#)
- compute_flux_gradient_layer, [5](#)
- compute_monthly_mean, [6](#)
- compute_neon_flux, [7](#)
- compute_surface_flux, [8](#)
- compute_surface_flux_layer, [9](#)
- correct_env_data, [9](#)
- dejong_shappert_flux, [10](#)
- depth_interpolate, [11](#)
- determine_position, [12](#)
- diffusivity, [12](#)
- env_fingerprint_plot, [13](#)
- fit_function, [14](#)
- flux_fingerprint_plot, [15](#)
- hirano_flux, [15](#)
- insert_mean, [16](#)
- inside_interval, [17](#)
- linear_regression, [18](#)
- loadByProduct, [3](#), [26](#)
- measurement_detect, [19](#)
- measurement_merge, [19](#)
- neonBoulded60Cal, [20](#)
- neonCal, [21](#)
- powerset, [22](#)
- quadrature_error, [23](#)
- reprocess_vswc, [24](#)
- sjer_env_data_2022_06, [24](#)
- sjer_flux_2022_06, [25](#)
- sjer_megapit_data_2022_06, [26](#)
- swc_correct, [27](#)
- swc_corrections, [28](#)
- tang_2003_flux, [28](#)
- tang_2005_flux, [29](#)