

# Package ‘rpls’

May 7, 2020

**Type** Package

**Title** Robust Partial Least Squares

**Version** 0.6.0

**Author** Peter Filzmoser, Sukru Acitas, Birdal Senoglu and Maximilian Plattner

**Maintainer** Peter Filzmoser <peter.filzmoser@tuwien.ac.at>

**Description** A robust Partial Least-Squares (PLS) method is implemented that is robust to outliers in the residuals as well as to leverage points. A specific weighting scheme is applied which avoids iterations, and leads to a highly efficient robust PLS estimator.

**License** GPL (>= 3)

**Imports** pcaPP, robustbase

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2020-05-07 13:30:02 UTC

## R topics documented:

pramml . . . . .	1
PRM . . . . .	3
ramml . . . . .	4

<b>Index</b>	<b>6</b>
--------------	----------

---

pramml *Partial Robust Adaptive Modified Maximum Likelihood*

---

## Description

Robust Adaptive Modified Maximum Likelihood (RAMML) estimators can be used in the context of PLS to obtain scores and loadings in the latent regression model. The corresponding method is called Partial RAMML (PRAMML).

**Usage**

```
pramml(X, y, a, reg = "lts", pmml, opt = "l1m", usesvd = FALSE)
```

**Arguments**

X	predictor matrix
y	response variable
a	number of PLS components
reg	regression procedure to be used to compute initial estimate of parameter for the linearization of the intractable term; choices are LTS regression ("lts") and S regression ("s")
pmml	shape parameter of long-tailed symmetric distribution (considered as robustness tuning constant)
opt	if "l1m" the mean centering is done by the l1-median; otherwise if "median" the coordinate-wise median is taken
usesvd	if TRUE singular value decomposition is performed; logical, default is FALSE

**Value**

coef	vector with regression coefficients
intercept	coefficient for intercept
wy	vector of length(y) with residual weights
wt	vector of length(y) with weights for leverage
w	overall weights
scores	matrix with PLS X-scores
loadings	matrix with PLS X-loadings
fitted.values	vector with fitted y-values
loadings	column means of X
fitted.values	mean of y

**Author(s)**

Sukru Acitas <sacitas@eskisehir.edu.tr>

**References**

S. Acitas, Robust Statistical Estimation Methods for High-Dimensional Data with Applications, tech. rep., TUBITAK 2219, International Post Doctoral Research Fellowship Programme, 2019.

**Examples**

```
U <- c(rep(2,20), rep(5,30))
X <- replicate(6, U+rnorm(50))
beta <- c(rep(1, 3), rep(-1,3))
e <- c(rnorm(45,0,1.5),rnorm(5,-20,1))
y <- X*%beta + e
res <- pramml(X, y, 4,"s", 16.5, opt ="l1m")
```

PRM

*Robust PLS***Description**

Robust PLS by partial robust M-regression.

**Usage**

```
PRM(formula,data,a,wfunX,wfunY,center.type,scale.type,usesvd,numit,prec)
```

**Arguments**

formula	an object of class formula
data	a data frame which contains the variables given in formula
a	number of PLS components
wfunX	weight function to downweight leverage points; predefined weight funcktions "Fair", "Huber", "Tukey" and "Hampel" with respective tuning constants are passed via a list object, e.g. list("Fair",0.95)
wfunY	weight function to downweight residuals; predefined weight funcktions "Fair", "Huber", "Tukey" and "Hampel" with respective tuning constants are passed via a list object, e.g. list("Fair",0.95)
center.type	type of centering of the data in form of a string that matches an R function, e.g. "median"
scale.type	type of scaling for the data in form of a string that matches an R function, e.g. "qn" or alternatively "no" for no scaling
numit	the number of maximal iterations for the convergence of the coefficient estimates
prec	a value for the precision of estimation of the coefficients
usesvd	if TRUE singular value decomposition is performed; logical, default is FALSE

**Details**

M regression is used to robustify PLS. Employment of seperate weight functions for leverage points and residuals.

**Value**

coef	vector with regression coefficients
intercept	coefficient for intercept
wy	vector of length(y) with residual weights
wt	vector of length(y) with weights for leverage
w	overall weights
scores	matrix with PLS X-scores
loadings	matrix with PLS X-loadings
fitted.values	vector with fitted y-values

**Author(s)**

Peter Filzmoser <peter.filzmoser@tuwien.ac.at>

**References**

S. Serneels, C. Croux, P. Filzmoser, and P.J. Van Espen. Partial robust M-regression. *Chemometrics and Intelligent Laboratory System*, Vol. 79(1-2), pp. 55-64, 2005.

**Examples**

```
U <- c(rep(2,20), rep(5,30))
X <- replicate(6, U+rnorm(50))
beta <- c(rep(1, 3), rep(-1,3))
e <- c(rnorm(45,0,1.5),rnorm(5,-20,1))
y <- X%%beta + e
d <- as.data.frame(X)
d$y <- y
res <- PRM(y~., data=d, 3, wfunX=list("Fair",0.95),
wfunY=list("Fair",0.95), center.type = "median",
scale.type = "no", usesvd = FALSE,
numit = 100, prec = 0.01)
res$coef
```

---

ramml

---

*Robust Adaptive Modified Maximum Likelihood*


---

**Description**

Modified Maximum Likelihood (MML) estimators are asymptotically equivalent to the ML estimators but their methodology works under the assumption of a known shape parameter. Robust Adaptive MML estimators weaken this assumption and are robust to vertical outliers as well as leverage points.

**Usage**

```
ramml(X,y,p,e)
```

**Arguments**

X	predictor matrix
y	response variable
p	shape parameter of long-tailed symmetric distribution (considered as robustness tuning constant)
e	parameter for the linearization of the intractable term

**Value**

coef	vector of coefficients
scale	estimate of sigma
fitted.values	vector with fitted y-values
residuals	vector with y-residuals

**Author(s)**

Sukru Acitas <sacitas@eskisehir.edu.tr>

**References**

S. Acitas, Robust Statistical Estimation Methods for High-Dimensional Data with Applications, tech. rep., TUBITAK 2219, International Post Doctoral Research Fellowship Programme, 2019.

# Index

\*Topic **multivariate**

pramml, 1

PRM, 3

ramml, 4

pramml, 1

PRM, 3

ramml, 4