

Package ‘PhysioIndexR’

November 12, 2025

Title Physiological and Stress Indices for Crop Evaluation

Version 0.1.0

Maintainer Vinayaka <vinayaka.b3vs@gmail.com>

Description

Crop production systems are increasingly challenged by climate variability, resource limitations, and biotic–abiotic stresses. In this context, stress tolerance indices and physiological trait estimators are essential tools to identify stable and superior genotypes, quantify yield stability under stress versus non-stress conditions, and understand plant adaptive responses. The 'PhysioIndexR' package provides a unified framework to compute commonly used stress indices, physiological traits, and derived metrics that are critical in crop improvement, crop physiology, and other agricultural sciences. The package includes functions to calculate classical stress tolerance indices (See Lamba et al., 2023; <[doi:10.1038/s41598-023-37634-8](https://doi.org/10.1038/s41598-023-37634-8)>) such as Tolerance (TOL), Stress Tolerance Index (STI), Stress Susceptibility Percentage Index (SSPI), Yield Index (YI), Yield Stability Index (YSI), Relative Stress Index (RSI), Mean Productivity (MP), Geometric Mean Productivity (GMP), Harmonic Mean (HM), Mean Relative Performance (MRP), and Percent Yield Reduction (PYR), along with a convenience wrapper `all_indices()` that returns all indices simultaneously. The function `mfvst_from_indices()` integrates these indices into a composite stress score using direction-aware membership values (0–1 scaling) and also averaging, facilitating genotype ranking and selection (See Vinu et al., 2025; <[doi:10.1007/s12355-025-01595-1](https://doi.org/10.1007/s12355-025-01595-1)>). The package also implements two novel composite functions: `WMFVST()`, which computes the Weighted Mean Membership Function Value for Stress Tolerance, and `WASI()`, which computes the Weighted Average Stress Index, both derived from membership function values (MFV) and raw stress index values, respectively. Beyond stress indices, the package provides functions for key physiological traits relevant to sugarcane and other crops: `bmap()` computes biomass accumulation and partitioning between leaf, cane/shoot, and root fractions. `chl()` estimates total chlorophyll content from Soil-Plant Analysis Development (SPAD) and Chlorophyll Content Index (CCI) values using validated quadratic models particularly for sugarcane (See Krishnapriya et al., 2020; <[doi:10.37580/JSR.2019.2.9.150-163](https://doi.org/10.37580/JSR.2019.2.9.150-163)>). `ctd()` calculates canopy temperature depression (CTD) from ambient and canopy temperatures, an important indicator of transpiration efficiency. `growth()` computes key growth analysis parameters, including Leaf Area Index (LAI), Net Assimilation Rate (NAR), and Crop Growth Rate (CGR) across crop growth stages (See Watson, 1958; <[doi:10.1093/oxfordjournals.aob.a083596](https://doi.org/10.1093/oxfordjournals.aob.a083596)>). `ranking()` provides flexible ranking utilities for genotype performance with multiple tie-handling and NA-placement op-

tions. Through these tools, the package enables researchers to: (i) quantify crop responses to stress environments, (ii) partition physiological components of yield, (iii) integrate multiple indices into composite metrics for genotype evaluation, and (iv) facilitate informed decision making in breeding pipelines, and plant physiology experiments. By combining physiology-based traits with quantitative stress indices, 'PhysioIndexR' supports comprehensive crop evaluation and helps researchers identify multi-stress-resilient superior genotypes, thereby contributing to genetic improvement and ensuring sustainable production of food, fuel, and fibre in the era of limited resources and climate change.

Imports stats

License GPL-3

Encoding UTF-8

Depends R (>= 4.1)

RoxygenNote 7.3.3

NeedsCompilation no

Author Vinayaka [aut, cre] (ORCID: <<https://orcid.org/0000-0001-5004-0084>>),
 Vengavasi Krishnapriya [aut, ctb] (ORCID:
 <<https://orcid.org/0000-0002-7496-5302>>),
 T. Lakshmi Pathy [aut, ctb] (ORCID:
 <<https://orcid.org/0000-0001-8940-7971>>),
 Amaresh [aut, ctb] (ORCID: <<https://orcid.org/0009-0000-5201-5755>>),
 K. Gopalareddy [aut, ctb] (ORCID:
 <<https://orcid.org/0000-0002-8825-6363>>),
 G.S. Suresha [aut, ctb] (ORCID:
 <<https://orcid.org/0000-0002-3283-6617>>),
 P. Govindaraj [aut, ctb]

Repository CRAN

Date/Publication 2025-11-12 10:50:01 UTC

Contents

all_indices	3
bmap	4
Chl	5
ctd	6
GMP	7
growth	8
HM	9
mfvst_from_indices	10
MP	11
MRP	12
PYR	13
ranking	14
RSI	15
SSPI	16
STI	17

<i>all_indices</i>	3
TOL	18
WASI	19
WMFVST	20
YI	21
YSI	22
Index	23

<i>all_indices</i>	<i>Computation of All Stress Indices at Once</i>
--------------------	--

Description

A convenience function that returns a data frame with 11 stress indices—Stress Tolerance (TOL), Stress Tolerance Index (STI), Stress Susceptibility Percentage Index (SSPI), Yield Index (YI), Yield Stability Index (YSI), Relative Stress Index (RSI), Mean Productivity (MP), Geometric Mean Productivity (GMP), Harmonic Mean (HM), Mean Relative Performance (MRP), and Percent Yield Reduction (PYR)—for the same given inputs (Lamba et al., 2023; doi:10.1038/s41598-023-37634-8).

Usage

```
all_indices(
  Gen,
  YN,
  YS,
  YMN = NULL,
  YMS = NULL,
  include_inputs = TRUE,
  name_vectors_by_gen = TRUE
)
```

Arguments

Gen	Character vector of Genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.
YMN, YMS	Optional numeric scalars for environment means. If NULL, computed.
include_inputs	Logical; if TRUE returns a list with YMN, YMS, and all.
name_vectors_by_gen	Ignored; kept for backward compatibility.

Value

If include_inputs=TRUE, a list with YMN, YMS, and all data frame.

References

Lamba, K., Kumar, M., Singh, V. et al. (2023). doi:10.1038/s41598-023-37634-8.

Examples

```
out <- all_indices(
  Gen=c("G1", "G2", "G3"),
  YN=c(10,8,5), YS=c(7,5,3)
)
print(out)
```

bmap

Biomass Accumulation and Partitioning (bmap)

Description

This function computes total above-ground biomass (TBM), root:shoot ratios, and biomass partitioning to leaf and cane/crop yield across stages/phases PI–PIII. For more details see Krishnapriya et al. (2020) [doi:10.37580/JSR.2019.2.9.150-163](https://doi.org/10.37580/JSR.2019.2.9.150-163). Here observations are recorded on the variables-Plant height (PH), Root length (RL), Leaf dry weight (LDW), Cane dry weight (CDW), and Root dry weight (RDW) at the distinct time interval (number of days between observations recorded) phases (PI-PIII)

Usage

```
bmap(data)
```

Arguments

data	<p>A data frame with column:</p> <ul style="list-style-type: none"> • Gen: Character vector of Genotype IDs. • LDW_PI, LDW_PII, LDW_PIII: Leaf dry weight (LDW) for three distinct phases I-III (PI-PIII). • CDW_PI, CDW_PII, CDW_PIII: Cane/crop dry weight (CDW) for three distinct phases I-III (PI-PIII). • RDW_PI: Root dry weight (RDW) for phase I (PI). • RL_PI: Root length (RL) for phase I (PI). • PH_PI: Plant height (PH) for phase I (PI).
------	--

Value

A list of result columns:

- Gen: Character vector of genotype IDs.
- TBM_PI, TBM_PII, TBM_PIII: Total above-ground biomass for three distinct phases I-III (PI-PIII).
- BPL_PI, BPL_PII, BPL_PIII: Biomass partitioning to leaf for three distinct phases I-III (PI-PIII).
- BPC_PI, BPC_PII, BPC_PIII: Biomass partitioning to cane (crop) for three distinct phases I-III (PI-PIII).
- RSRDW: Root-to-shoot ratio based on dry weight
- RSRL: Root-to-shoot ratio based on length

References

Krishnapriya, V., Arunkumar, R., Gomathi, R. and Vasantha, S. (2020). doi:[10.37580/JSR.2019.2.9.150-163](https://doi.org/10.37580/JSR.2019.2.9.150-163).

Examples

```
# Creating a sample dataset
df <- data.frame(
  Gen = c("V1", "V2", "V3"),
  PH_PI = c(161.0, 144.0, 158.0),
  PH_PII = c(212.0, 189.0, 200.0),
  PH_PIII = c(243.0, 223.0, 240.0),
  RL_PI = c(98.2, 83.5, 72.7),
  LDW_PI = c(83.9, 136.1, 137.6),
  CDW_PI = c(36.1, 55.9, 57.3),
  RDW_PI = c(17.7, 19.2, 12.4),
  LDW_PII = c(138.5, 193.2, 183.0),
  CDW_PII = c(76.2, 136.4, 145.0),
  LDW_PIII = c(292.2, 386.5, 450.1),
  CDW_PIII = c(190.4, 336.8, 323.4)
)
bmap(df)
```

Chl

Chlorophyll concentration from non-destructive indices (Chl)

Description

This function computes total chlorophyll content derived from SPAD (Soil-Plant Analysis Development) value (TCHLSPAD), and total chlorophyll content derived from CCI (Chlorophyll Content Index) value (TCHLCCI). For more details see Krishnapriya et al. (2020) doi:[10.37580/JSR.2019.2.9.150-163](https://doi.org/10.37580/JSR.2019.2.9.150-163).

Usage

```
Chl(Gen, SPAD_PI, CCI_PI)
```

Arguments

Gen	Character vector of Genotype IDs.
SPAD_PI	Numeric vector of SPAD (Soil-Plant Analysis Development) values for phase I (PI).
CCI_PI	Numeric vector of CCI (Chlorophyll Content Index) values for phase I (PI).

Value

A list of output columns:

- Gen: Character vector of genotype IDs.
- TCHLSPAD: total chlorophyll content derived from SPAD (Soil-Plant Analysis Development) value (TCHLSPAD).
- TCHLCCI: total chlorophyll content derived from CCI (Chlorophyll Content Index) value (TCHLCCI).

References

Krishnapriya, V., Arunkumar, R., Gomathi, R. and Vasantha, S. (2020). doi:[10.37580/JSR.2019.2.9.150-163](https://doi.org/10.37580/JSR.2019.2.9.150-163).

Examples

```
# Creating a sample dataset
Gen = c("V1", "V2", "V3")
SPAD_PI = c(43.1, 44.6, 38.6)
CCI_PI = c(27.3, 30.4, 21.3)
Chl(Gen, SPAD_PI, CCI_PI)
```

ctd

Canopy Temperature Depression (CTD)

Description

This function computes canopy temperature depression (CTD). For more details see Watson (1958) doi:[10.1093/oxfordjournals.aob.a083596](https://doi.org/10.1093/oxfordjournals.aob.a083596).

Usage

```
ctd(Gen, amb.temp, CT_PI)
```

Arguments

Gen	Character vector of genotype IDs.
amb.temp	Numeric scalar; ambient temperature (°C) (user must define this input).
CT_PI	Numeric vector; canopy temperature (°C) at phase I (PI).

Value

A list of output columns:

- Gen: Character vector of genotype IDs.
- CTD: Canopy temperature depression (CTD) values for respective genotypes.

References

Watson, D.J. (1958). doi:[10.1093/oxfordjournals.aob.a083596](https://doi.org/10.1093/oxfordjournals.aob.a083596).

Examples

```
# Creating a sample dataset
df <- data.frame(
  Gen = c("V1", "V2", "V3"),
  CT_PI = c(26.7, 31.4, 28.4)
)
ctd(df$Gen, amb.temp = 34.2, CT_PI = df$CT_PI)
```

GMP

Geometric Mean Productivity (GMP)

Description

This function computes Geometric Mean Productivity (GMP) using any traits (like yield) under stress and non-stress conditions. The lower values of GMP indicates greater tolerance. For more details see Fernandez (1992).

Usage

```
GMP(Gen, YN, YS)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.

Value

A data frame with Gen, GMP.

$$GMP = \sqrt{YN \times YS}$$

References

Fernandez, G.C.J. (1992). Effective selection criteria for assessing plant stress tolerance. In: Proceedings of the international symposium on adaptation of vegetables and other food crops in temperature and water stress. AVRDC Publication: Tainan, Taiwan: Shanhua: Chapter (25), 257–270.

Examples

```

out=GMP(
  Gen=c("G1", "G2", "G3"),
  YN=c(10,8,5),
  YS=c(7,5,3)
)
print(out)

```

growth

Growth Indices

Description

This function computes genotype (Gen) wise growth indices-leaf area index (LAI), net assimilation rate (NAR), and crop growth rate (CGR) for time intervals between distinct stages/phases (days). For more details see Williams (1946), and Watson (1958) [doi:10.1093/oxfordjournals.aob.a083596](https://doi.org/10.1093/oxfordjournals.aob.a083596).

Usage

```
growth(data, gr.area, t.interval1, t.interval2)
```

Arguments

data	Data frame with input columns: <ul style="list-style-type: none"> • Gen: Character vector of Genotype IDs. • LDW_PI, LDW_PII, LDW_PIII: Leaf dry weight (LDW) for three distinct phases I-III (PI-PIII). • CDW_PI, CDW_PII, CDW_PIII: Cane/crop dry weight (CDW) for three distinct phases I-III (PI-PIII). • LA_PI, LA_PII, LA_PIII): Leaf area (LA) for three distinct phases I-III (PI-PIII).
gr.area	Ground area occupied by the sample (cm ² or m ² , same unit as that of leaf area) (user must define this input).
t.interval1	Time interval (days) between consecutive sampling phases PI and PII (user must define this input).
t.interval2	Time interval (days) between consecutive sampling phases PII and PIII (user must define this input).

Value

A list of result components:

- Gen: Character vector of genotype IDs.
- LAI_PI, LAI_PII, LAI_PIII): Leaf Area Index (LAI) for three distinct phases I-III (PI-PIII).

- NAR_PII, NAR_PIII: Net Assimilation Rate (NAR) for two consecutive sampling intervals (PI–PII) and (PII–PIII).
- CGR_PII, CGR_PIII: Crop Growth Rate (CGR) for two consecutive sampling intervals (PI–PII) and (PII–PIII).

References

Williams, R.F. (1946). The physiology of plant growth with special reference to the concept of net assimilation rate. *Annals of Botany*, 10(37), 41-72.

Watson, D.J. (1958). doi:[10.1093/oxfordjournals.aob.a083596](https://doi.org/10.1093/oxfordjournals.aob.a083596).

Examples

```
# Creating a sample dataset
df <- data.frame(
  Gen = c("V1", "V2", "V3"),
  LA_PI = c(599.4, 544.4, 573.2),
  LA_PII = c(1533.4, 1088.0, 1633.1),
  LA_PIII= c(1111.2, 866.0, 1181.0),
  LDW_PI = c(83.9, 136.1, 137.6),
  CDW_PI = c(36.1, 55.9, 57.3),
  LDW_PII= c(138.5, 193.2, 183.0),
  CDW_PII= c(76.2, 136.4, 145.0),
  LDW_PIII=c(292.2, 386.5, 450.1),
  CDW_PIII=c(190.4, 336.8, 323.4)
)
growth(df, gr.area = 250, t.interval1 = 30, t.interval2 = 60)
```

HM

Harmonic Mean (HM)

Description

This function computes Harmonic Mean (HM) using any traits (like yield) under stress and non-stress conditions. The lower values of HM indicates greater tolerance. For more details see Bidinger et al. (1987) doi:[10.1071/AR9870037](https://doi.org/10.1071/AR9870037).

Usage

```
HM(Gen, YN, YS)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.

Value

A data frame with Gen, HM.

$$HM = \frac{2 \times YN \times YS}{YN + YS}$$

References

Bidinger, F.R., Mahalakshmi, V. and Rao, G.D.P. (1987). doi:[10.1071/AR9870037](https://doi.org/10.1071/AR9870037).

Examples

```
out = HM(
  Gen=c("G1", "G2", "G3"),
  YN=c(10, 8, 5),
  YS=c(7, 5, 3)
)
print(out)
```

mfvst_from_indices *Membership Function Value for Stress Tolerance (MFVST)*

Description

This function computes membership function scores (0..1) for each available index column using min-max scaling with direction awareness, then aggregates them into a simple average MFVST. For more details see Vinu et al. (2025) doi:[10.1007/s12355-025-01595-1](https://doi.org/10.1007/s12355-025-01595-1).

Usage

```
mfvst_from_indices(
  df,
  gen_col = "Gen",
  lower_better = c("TOL", "SSPI", "RSI", "PYR"),
  higher_better = c("STI", "YI", "YSI", "MP", "GMP", "HM", "MRP"),
  weights = NULL,
  robust = FALSE,
  probs = c(0.01, 0.99)
)
```

Arguments

df	A data frame containing the stress indices (e.g., from <code>all_indices()</code> \$all).
gen_col	Name of genotype column; if present, it is included in the output.
lower_better	Character vector listing indices where a lower value is preferred.
higher_better	Character vector listing indices where a higher value is preferred.

weights	Optional named numeric vector of weights for indices.
robust	Logical; if TRUE, winsorizes by probs.
probs	Two-element numeric vector of quantiles for robust winsorization.

Value

A list with \$MFVST_indexwise: a data frame that contains per-index membership columns (suffix "_M") and the average MFVST, that is, Mean_MFVST.

References

Vinu, V., Lakshmi Pathy, T., Mahadeva Swamy, H.K. et al. (2025). doi:10.1007/s12355-025-01595-1.

Examples

```
df <- all_indices(
  Gen=c("G1", "G2", "G3"),
  YN=c(10, 8, 5),
  YS=c(7, 5, 3)
)
df1 <- as.data.frame(df$all)
mfvst <- mfvst_from_indices(df1)
print(mfvst)
```

MP	<i>Mean Productivity (MP)</i>
----	-------------------------------

Description

This function computes Mean Productivity (MP) using any traits (like yield) under stress and non-stress conditions. The lower values of MP indicates greater tolerance. For more details see Rosielle and Hamblin (1981) doi:10.2135/cropsci1981.0011183X002100060033x.

Usage

```
MP(Gen, YN, YS)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.

Value

A data frame with Gen, MP.

$$MP = \frac{YN + YS}{2}$$

References

Rosielle, A.A. and Hamblin, J. (1981). <10.2135/cropsci1981.0011183X002100060033x>.

Examples

```
out = MP(
  Gen=c("G1", "G2", "G3"),
  YN=c(10, 8, 5),
  YS=c(7, 5, 3)
)
print(out)
```

 MRP

Mean Relative Performance (MRP)

Description

This function computes Mean Relative Performance (MRP) using any traits (like yield) under stress and non-stress conditions. The lower values of MRP indicates greater tolerance. For more details see Ramirez-Vallejo and Kelly (1998) [doi:10.1023/A:1018353200015](https://doi.org/10.1023/A:1018353200015).

Usage

```
MRP(Gen, YN, YS, YMN = NULL, YMS = NULL)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.
YMN	Optional numeric scalar: mean of YN. If NULL, computed.
YMS	Optional numeric scalar: mean of YS. If NULL, computed.

Value

A list with YMN, YMS, and Result (data frame with Gen, MRP).

$$MRP = \left(\frac{YS}{YMS} \right) + \left(\frac{YN}{YMN} \right)$$

References

Ramirez-Vallejo, P. and Kelly, J.D. (1998). [doi:10.1023/A:1018353200015](https://doi.org/10.1023/A:1018353200015).

Examples

```

out = MRP(
  Gen=c("G1", "G2", "G3"),
  YN=c(10, 8, 5),
  YS=c(7, 5, 3)
)
print(out)

```

 PYR

Percent Yield Reduction (PYR)

Description

This function computes Percent Yield Reduction (PYR) using any traits (like yield) under stress and non-stress conditions. The lower values of PYR indicates greater tolerance. For more details see Farshadfar and Javadinia (2011).

Usage

```
PYR(Gen, YN, YS)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.

Value

A data frame with Gen, PYR.

$$PYR = \frac{(YN - YS)}{YN} \times 100$$

References

Farshadfar, E. and Javadinia, J. (2011). Evaluation of chickpea (*Cicer arietinum* L.) genotypes for drought tolerance. *Seed and Plant Improvement Journal*, 27(4), 517–537.

Examples

```

out = PYR(
  Gen=c("G1", "G2", "G3"),
  YN=c(10, 8, 5),
  YS=c(7, 5, 3)
)
print(out)

```

ranking

Flexible Ranking Utility

Description

A thin wrapper around base `rank` with support for ascending/descending order, multiple tie strategies, NA placement, and dense ranks.

Usage

```
ranking(  
  v,  
  direction = c("asc", "desc"),  
  ties = c("average", "min", "max", "first", "random"),  
  na.last = c("keep", "bottom", "top"),  
  dense = FALSE  
)
```

Arguments

<code>v</code>	Numeric (or coercible) vector to rank.
<code>direction</code>	Character, one of "asc" or "desc".
<code>ties</code>	Character, tie-breaking: one of "average", "min", "max", "first", "random".
<code>na.last</code>	Character, placement of NAs: "keep", "bottom", or "top".
<code>dense</code>	Logical; if TRUE, returns dense ranks (1,2,3,...) without gaps.

Value

An integer/numeric vector of ranks, same length as `v`.

Examples

```
ranking(c(3, 3, 2, NA, 5), direction="asc", ties="min", na.last="bottom")  
ranking(c(3, 3, 2, 5), direction = "desc", dense = TRUE)  
Gen=c("G1", "G2", "G3")  
YN=c(10,8,5)  
YS=c(7,5,3)  
a=STI(Gen, YN, YS) # for instance STI taken here.  
out=data.frame(a$Result$Gen, a$Result$STI,  
  ranking(a$Result$STI, direction="desc")  
)  
print(out)
```

RSI

*Relative Stress Index (RSI)***Description**

This function computes Relative Stress Index (RSI) using any traits (like yield) under stress and non-stress conditions. The lower values of RSI indicates greater tolerance. For more details see Fischer and Wood (1979) [doi:10.1071/AR9791001](https://doi.org/10.1071/AR9791001).

Usage

```
RSI(Gen, YN, YS, YMN = NULL, YMS = NULL)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.
YMN	Optional numeric scalar: mean of YN. If NULL, computed.
YMS	Optional numeric scalar: mean of YS. If NULL, computed.

Value

A list with YMN, YMS, and Result (data frame with Gen, RSI).

$$RSI = \frac{(YN/YS)}{(YMS/YMN)}$$

References

Fischer, R.A. and Wood, J.T. (1979). Drought resistance in spring wheat cultivars. III.* Yield associations with morpho-physiological traits. Australian Journal of Agricultural Research, 30(6), 1001-1020.

Examples

```
out = RSI(
  Gen=c("G1", "G2", "G3"),
  YN=c(10, 8, 5),
  YS=c(7, 5, 3)
)
print(out)
```

 SSPI

Stress Susceptibility Percentage Index (SSPI)

Description

This function computes Stress Susceptibility Percentage Index (SSPI) using any traits (like yield) under stress and non-stress conditions. The lower values of SSPI indicates greater tolerance. For more details see Moosavi et al. (2008).

Usage

```
SSPI(Gen, YN, YS, YMN = NULL)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.
YMN	Optional numeric scalar: mean of YN. If NULL, computed.

Value

A list with YMN and Result (data frame with Gen, SSPI).

$$SSPI = \frac{(YN - YS)}{2 \times YMN} \times 100$$

References

Mousavi, S.S., YAZDI, S.B., Naghavi, M.R., Zali, A.A., Dashti, H. and Pourshahbazi, A. (2008). Introduction of new indices to identify relative drought tolerance and resistance in wheat genotypes. Desert 12, 165–178.

Examples

```
out=SSPI(Gen=c("G1","G2","G3"), YN=c(10,8,5), YS=c(7,5,3))
print(out)
```

 STI *Stress Tolerance Index (STI)*

Description

This function computes stress tolerance index (STI) using any traits (like yield) under stress and non-stress conditions. The lower values of STI indicates greater tolerance. For more details see Fernandez (1992).

Usage

```
STI(Gen, YN, YS, YMN = NULL)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.
YMN	Optional numeric scalar: mean of YN. If NULL, computed as <code>mean(YN, na.rm=TRUE)</code> .

Value

A list of components:

- `YMN`: Mean of yield (any trait) values under normal condition
- `Result`: It includes
 - `Gen`: Character vector of genotype IDs.
 - `STI`: Estimated stress tolerance index (STI) values for respective genotypes.

$$STI = \frac{YN \times YS}{(YMN)^2}$$

References

Fernandez, G.C.J. (1992). Effective selection criteria for assessing plant stress tolerance. In: Proceedings of the international symposium on adaptation of vegetables and other food crops in temperature and water stress. AVRDC Publication: Tainan, Taiwan: Shanhua: Chapter (25), 257–270.

Examples

```
out=STI(Gen=c("G1", "G2", "G3"), YN=c(10, 8, 5), YS=c(7, 5, 3))
print(out)
```

TOL	<i>Stress Tolerance (TOL)</i>
-----	-------------------------------

Description

This function computes stress tolerance (TOL) using any traits like yield under stress and non-stress conditions. The lower values of TOL indicates greater tolerance. For more details see Rosielle and Hamblin (1981) [doi:10.2135/cropsci1981.0011183X002100060033x](https://doi.org/10.2135/cropsci1981.0011183X002100060033x).

Usage

```
TOL(Gen, YN, YS)
```

Arguments

Gen	Character vector of Genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.

Value

A list of output columns:

- Gen: Character vector of genotype IDs.
- TOL: Estimated stress tolerance (TOL) values for respective genotypes.

$$TOL = YN - YS$$

References

Rosielle, A.A. and Hamblin, J. (1981). [doi:10.2135/cropsci1981.0011183X002100060033x](https://doi.org/10.2135/cropsci1981.0011183X002100060033x).

Examples

```
TOL(Gen=c("G1", "G2", "G3"), YN=c(10, 8, 5), YS=c(7, 5, 3))
```

WASI

Weighted Average Stress Index (WASI)

Description

A composite measure that computes the weighted mean for each genotype across multiple stress indices, accounting for whether higher or lower values are better.

Usage

```
WASI(data, decimals = 5)
```

Arguments

<code>data</code>	A data frame containing genotype IDs (Gen) and stress index values (GMP, HM, MP, MRP, PYR, RSI, SSPI, STI, TOL, YI, YSI).
<code>decimals</code>	Integer; number of decimal places to use for Excel-style data. All indices are rounded to this precision before ranking and again for the index × rank products (default 5).

Details

The Weighted Average Stress Index for genotype i is

$$WASI_i = \frac{\sum_{j=1}^k (X_{ij} \times R_{ij})}{\sum_{j=1}^k R_{ij}},$$

where X_{ij} is the value of genotype i for index j , and R_{ij} is its rank, determined by whether higher or lower values are favorable. Ranks use Excel-like behavior (`ties.method = "min"`). Indices are rounded to `decimals` places prior to ranking and multiplication to better match Excel-style calculations.

Value

A data frame with Gen and its computed WASI (sorted in descending order).

Examples

```
df <- data.frame(
  Gen = paste0("G", 1:5),
  GMP = c(1:5),
  HM = c(6:10),
  MP = c(11:15),
  MRP = c(16:20),
  PYR = c(21:25),
  RSI = c(26:30),
  SSPI = c(31:35),
  STI = c(0.1, 0.2, 0.3, 0.4, 0.5),
```

```
TOL = c(41:45),
YI = c(0.6, 0.7, 0.8, 0.9, 1.0),
YSI = c(0.2, 0.3, 0.4, 0.5, 0.6)
)
WASI(df)
```

WMFVST	<i>Weighted Mean Membership Function Value for Stress Tolerance (WMFVST)</i>
--------	--

Description

Its a composite measure which computes the weighted mean of MFVST values (0–1 range) for each genotype across multiple stress indices, considering whether higher or lower values are better.

Usage

```
WMFVST(data)
```

Arguments

data	A data frame containing genotype IDs (Gen) and MFVST values for the indices (GMP, HM, MP, MRP, PYR, RSI, SSPI, STI, TOL, YI, YSI), each within the range of 0 to 1.
------	---

Details

The Weighted Mean MFVST for genotype i is:

$$WMFVST_i = \frac{\sum_{j=1}^k (MFVST_{ij} \times R_{ij})}{\sum_{j=1}^k R_{ij}}$$

where $MFVST_{ij}$ is the MFVST value of genotype i for index j , and R_{ij} is its rank, determined by whether higher or lower values are favorable.

Value

A data frame with Gen and its Weighted Mean MFVST (WMFVST).

Examples

```
set.seed(123)
df <- data.frame(
  Gen = paste0("G", 1:5),
  GMP = runif(5),
  HM = runif(5),
  MP = runif(5),
  MRP = runif(5),
```

```

PYR = runif(5),
RSI = runif(5),
SSPI = runif(5),
STI = runif(5),
TOL = runif(5),
YI = runif(5),
YSI = runif(5)
)
WMFVST(df)

```

YI *Yield Index (YI)*

Description

This function computes Yield Index (YI) using any traits (like yield) under stress and non-stress conditions. The lower values of YI indicates greater tolerance. For more details see Gavuzzi et al. (1997) [doi:10.4141/P96-130](https://doi.org/10.4141/P96-130).

Usage

```
YI(Gen, YS, YMS = NULL)
```

Arguments

Gen	Character vector of genotype identifiers.
YS	Numeric vector: yield (any trait) under stress environment.
YMS	Optional numeric scalar: mean of YS. If NULL, computed.

Value

A list with YMS and Result (data frame with Gen, YI).

$$YI = \frac{YS}{YMS}$$

References

Gavuzzi, P., Rizza, F., Palumbo, M., Campanile, R.G., Ricciardi, G.L. and Borghi, B. (1997). Evaluation of field and laboratory predictors of drought and heat tolerance in winter cereals. *Canadian Journal of Plant Science*, 77(4), 523-531.

Examples

```

out = YI(Gen=c("G1", "G2", "G3"), YS=c(7, 5, 3))
print(out)

```

YSI

Yield Stability Index (YSI)

Description

This function computes Yield Stability Index (YSI) using any traits (like yield) under stress and non-stress conditions. The lower values of YSI indicates greater tolerance. For more details see Bouslama and Schapaugh (1984) [doi:10.2135/cropsci1984.0011183X002400050026x](https://doi.org/10.2135/cropsci1984.0011183X002400050026x).

Usage

```
YSI(Gen, YN, YS)
```

Arguments

Gen	Character vector of genotype identifiers.
YN	Numeric vector: yield (any trait) under non-stress (normal) environment.
YS	Numeric vector: yield (any trait) under stress environment.

Value

A data frame with columns Gen, YSI.

$$YSI = \frac{YS}{YN}$$

References

Bouslama, M. and Schapaugh Jr, W.T. (1984). [doi:10.2135/cropsci1984.0011183X002400050026x](https://doi.org/10.2135/cropsci1984.0011183X002400050026x).

Examples

```
out = YSI(Gen=c("G1", "G2", "G3"), YN=c(10, 8, 5), YS=c(7, 5, 3))
print(out)
```

Index

[all_indices](#), 3

[bmap](#), 4

[Ch1](#), 5

[ctd](#), 6

[GMP](#), 7

[growth](#), 8

[HM](#), 9

[mfvst_from_indices](#), 10

[MP](#), 11

[MRP](#), 12

[PYR](#), 13

[rank](#), 14

[ranking](#), 14

[RSI](#), 15

[SSPI](#), 16

[STI](#), 17

[TOL](#), 18

[WASI](#), 19

[WMFVST](#), 20

[YI](#), 21

[YSI](#), 22